

BROMIKA[®] ELIXIR

SECRETOLYTIC AGENT COUGH SYRUP

COMPOSITION :

Each 5 ml elixir contains :

Bromhexine HCl 4 mg
Ethanol 3,5 %

MODE OF ACTION :

Bromika reduces the viscosity of respiratory tract secretions by changing the structure of mucoproteins and mucopolysaccharides in sputum and thereby easing difficulty in expectoration.

INDICATIONS :

Bromika is indicated for the treatment of productive coughs to facilitate expectoration, such as cough due to influenza or respiratory tract inflammation.

DOSAGE :

Children 2 - 5 years : 2 x 1 measuring spoonful.
Children 5 - 10 years : 3 x 1 measuring spoonful.
Adults & Children > 10 years : 3 x 2 measuring spoonful.
Or according to the physician.

WARNING AND PRECAUTIONS :

Bromika should be given cautiously to patients with gastric ulceration and to pregnant women during the first trimester of pregnancy and lactating women.

ADVERSE EFFECTS :

The following adverse effects may occur : nausea, diarrhea, gastrointestinal disturbances, but these are almost invariably mild. Headache, vertigo, sweating, skin rash and a transient rise in transaminase have been reported.

DRUG INTERACTION :

Concomitant administration of antibiotics (amoxicillin, cefuroxime, doxycycline) will increase the antibiotic concentration.

PRESENTATION :

Bottles containing 60 ml, 120 ml. Reg. No. : DTL 9709312434 A1

STORE IN A COOL AND DRY PLACE

Manufactured by :



PT. IKAPHARMINDO PUTRAMAS
PHARMACEUTICAL LABORATORIES
JAKARTA - INDONESIA